

# Novartis Gains FDA Approval for Afinitor® as First New Treatment in Nearly Three Decades for Patients with Advanced Pancreatic NET

May 07, 2011

- - Data show Afinitor delays tumor growth and reduces risk of disease progression in patients with advanced neuroendocrine tumors (NET) of pancreatic origin(1)
- - Afinitor represents a new approach to treat advanced pancreatic NET, an aggressive cancer for which there has been limited treatment options(1),(2),(3)

EAST HANOVER, N.J., May 5, 2011 /PRNewswire/ -- Novartis Pharmaceuticals Corporation ("Novartis") announced today that the US Food and Drug Administration (FDA) approved Afinitor® (everolimus) tablets for the treatment of progressive neuroendocrine tumors of pancreatic origin (PNET) in patients with unresectable, locally advanced or metastatic disease(4). This marks the first approval of a treatment for this patient population in the US in nearly 30 years(5).

The approval was based on Phase III data from the RADIANT-3 (RAD001 In Advanced Neuroendocrine Tumors) trial, showing treatment with Afinitor more than doubled the time without tumor growth (median 4.6 to 11.0 months) and reduced the risk of cancer progression by 65% when compared with placebo in patients with advanced pancreatic NET (hazard ratio=0.35 [95% confidence interval (CI), 0.27 to 0.45]; p<0.001). A consistent improvement in progression-free survival was seen with Afinitor in all patient subgroups(1). The FDA determined that the safety and effectiveness of Afinitor in the treatment of patients with carcinoid tumors have not been established(4).

"The FDA approval of Afinitor represents an important step forward for patients with advanced pancreatic NET," said James Yao, MD, Associate Professor of Medicine, The University of Texas MD Anderson Cancer Center, Houston, Texas. "Patients will now have access to a treatment that has been shown to significantly delay tumor growth and reduce the risk of disease progression."

Approximately 60% of pancreatic NET patients are diagnosed with advanced disease(2). This means that the cancer has already spread to other parts of the body, and is considered aggressive and difficult to treat(3). The five-year survival rate for these patients is 27%(6).

"With this approval, physicians can now offer their patients with progressive pancreatic NET a new treatment helping to fulfill a critical unmet need," said Herve Hoppenot, President, Novartis Oncology. "This is the third indication for Afinitor in the US in just over two years, providing further evidence that inhibiting mTOR plays an important role in treating multiple tumor types."

Afinitor targets mTOR, a protein that acts as an important regulator of tumor cell division, blood vessel growth and cell metabolism(7). Preclinical and clinical data have established the role of mTOR in the development and progression of several types of tumors, including advanced pancreatic NET(1),(7).

About neuroendocrine tumors of pancreatic origin (pancreatic NET)

Neuroendocrine tumors arise from cells that can produce and secrete a variety of hormones that regulate

bodily functions(8). These tumors can occur anywhere in the body; however, most are found in the pancreas (pancreatic NET), gastrointestinal tract or lungs (carcinoid tumors)(6),(9). Pancreatic NET, also known as islet cell tumors, is a rare type of cancer different from pancreatic exocrine cancer, which is generally referred to as pancreatic cancer(3),(10). There have been limited treatment options for patients with pancreatic NET(2).

### About RADIANT-3

RADIANT-3 is a Phase III prospective, double-blind, randomized, parallel group, placebo-controlled, multicenter study. The trial examined the efficacy and safety of Afinitor plus best supportive care (BSC) versus placebo plus BSC in 410 patients with advanced, low- or intermediate-grade pancreatic NET. Patients who met the study entry criteria were randomized 1:1 to receive either Afinitor 10 mg once-daily (n=207) or daily placebo (n=203) orally, both in conjunction with BSC(1).

The primary endpoint of RADIANT-3 is progression-free survival. Secondary endpoints include safety, objective response rate (confirmed according to RECIST), duration of response and overall survival(1).

### About Afinitor (everolimus)

Afinitor® (everolimus) tablets is approved in the US for the treatment of progressive neuroendocrine tumors of pancreatic origin (PNET) in patients with unresectable, locally advanced or metastatic disease. The safety and effectiveness of Afinitor in the treatment of patients with carcinoid tumors have not been established.

For more information visit [www.AFINITOR.com/pNET](http://www.AFINITOR.com/pNET) or call 1-888-4-AFINITOR. US patients who may be eligible for financial assistance can learn about the Patient Assistance Now Oncology (PANO) reimbursement support program by contacting 1-800-282-7630 or visiting the Afinitor website.

Afinitor is approved in the US for the treatment of patients with advanced renal cell carcinoma (RCC) after failure of treatment with sunitinib or sorafenib and in the European Union (EU) for the treatment of patients with advanced RCC whose disease has progressed on or after treatment with vascular endothelial growth factor (VEGF)-targeted therapy.

Afinitor is also approved in the US to treat patients with subependymal giant cell astrocytoma (SEGA) associated with tuberous sclerosis who require therapeutic intervention but are not candidates for curative surgical resection. The effectiveness of Afinitor is based on an analysis of change in SEGA volume. Clinical benefit such as improvement in disease-related symptoms or increase in overall survival has not been shown. Novartis has submitted marketing applications for everolimus for this use to the European Medicines Agency (EMA) and the Swiss Agency for Therapeutic Products (Swissmedic), and additional regulatory submissions are under way worldwide.

Afinitor is available in the US in 2.5 mg, 5 mg and 10 mg tablet strengths.

In the US, everolimus is available in different dosage strengths for the non-oncology patient population under the trade name Zortress® for the prophylaxis of organ rejection in adult patients at low-moderate immunologic risk receiving a kidney transplant. In the EU, everolimus is available in different dosage strengths under the trade name Certican® for the prevention of organ rejection in heart and kidney transplant recipients.

Everolimus is exclusively licensed to Abbott and sublicensed to Boston Scientific for use in drug-eluting stents.

Not all indications are available in every country. As an investigational compound the safety and efficacy profile of everolimus has not yet been established in all countries in pancreatic or any other type of NET. Access to everolimus outside of the approved indications has been carefully controlled and monitored in clinical trials designed to better understand the potential benefits and risks of the compound. Because of the

uncertainty of clinical trials, there is no guarantee that everolimus will become commercially available for pancreatic or any other type of NET, or additional indications anywhere else in the world.

#### Important Safety Information for Afinitor® (everolimus) tablets

Patients should not take Afinitor if they are allergic to Afinitor or to any of its ingredients. Patients should tell their healthcare provider before taking Afinitor if they are allergic to sirolimus (Rapamune®) or temsirolimus (Torisel®).

Afinitor can cause serious side effects including lung or breathing problems, infections and kidney failure, which can lead to death. If patients experience these serious side effects, they may need to stop taking Afinitor for a while or use a lower dose. Patients should tell their healthcare provider right away if they have any of these symptoms: new or worsening cough, shortness of breath, chest pain, difficulty breathing or wheezing.

Afinitor may make patients more likely to develop an infection, such as pneumonia, or a bacterial, fungal or viral infection. Viral infections may include reactivation of hepatitis B in people who have had hepatitis B in the past. In some people these infections may be severe, and can even lead to death. Patients may need to be treated as soon as possible. Patients should tell their healthcare provider right away if they have a temperature of 100.5 degrees F or above, chills or do not feel well. Symptoms of hepatitis B or infection may include the following: fever, skin rash, joint pain and inflammation, tiredness, loss of appetite, nausea, pale stool or dark urine, yellowing of the skin or pain in the upper right side.

Afinitor can cause mouth ulcers and sores, which are the most frequently occurring side effects occurring in approximately 44%-64% advanced kidney cancer and advanced pancreatic NET patients taking Afinitor. Eighty-six percent of patients taking Afinitor for SEGA developed mouth ulcers/sores. Patients should tell their healthcare provider if they have pain, discomfort or open sores in their mouth. Their healthcare provider may tell them to use a special mouthwash or mouth gel that does not contain alcohol or peroxide.

Afinitor may cause kidney failure. In some people this may be severe and can even lead to death. Patients should have tests to check their kidney function before and during their treatment with Afinitor.

Patients will have regular blood tests before they start and as needed during their treatment with Afinitor. These tests will include tests to check the patient's blood cell count, kidney and liver function and blood sugar levels. Patients who receive Afinitor for the treatment of SEGA will need regular blood tests to measure how much Afinitor is in their blood since this will help their doctor decide how much Afinitor they need to take.

Afinitor may affect the way other medicines work, and other medicines can affect how Afinitor works. Using Afinitor with other medicines can cause serious side effects. Patients should tell their healthcare provider about all of the medicines they take, including prescription and non-prescription medicines, vitamins and herbal supplements such as: St. John's Wort, and medicine for fungal infections, bacterial infections, tuberculosis, seizures, HIV-AIDS, heart conditions or high blood pressure and medicines that suppress their immune system. Patients should not drink grapefruit juice or eat grapefruit during their treatment with Afinitor as it may make the amount of Afinitor in their blood increase to a harmful level.

Patients should not take Afinitor tablets which are broken or crushed. Patients should not chew or crush the tablets.

The amount of Afinitor in the blood was increased in patients who had liver problems. Patients should tell their healthcare provider about all their medical conditions, including if they have or have had liver problems, diabetes or high blood sugar, high cholesterol levels, infections, hepatitis B or other medical conditions.

Patients should tell their healthcare provider if they are scheduled to receive any vaccinations. Patients should

not receive a live vaccine or be around people who have recently received a live vaccine during treatment with Afinitor.

It is not known if Afinitor will harm an unborn baby. Women should use effective birth control while using Afinitor and for eight weeks after stopping treatment.

Common side effects of Afinitor in patients with advanced pancreatic neuroendocrine tumors include mouth ulcers, rash, diarrhea, swelling of arms, hands, feet, ankles, face or other parts of the body, abdominal pain, nausea, fever and headache. Common side effects of Afinitor in patients with advanced kidney cancer include mouth ulcers, infections, feeling weak or tired, cough and diarrhea. Common side effects of Afinitor in patients with SEGA include mouth ulcers, infections of the respiratory tract, sinuses and ears and fever.

Please see full [Prescribing Information](#) for Afinitor.

#### Disclaimer

The foregoing release contains forward-looking statements that can be identified by terminology such as "will," "risk," "under way," "potential," or similar expressions, or by express or implied discussions regarding potential submissions or approvals for new indications or labeling for Afinitor, or regarding the potential timing of any such submissions or approvals, or regarding potential future revenues from Afinitor. You should not place undue reliance on these statements. Such forward-looking statements reflect the current views of management regarding future events, and involve known and unknown risks, uncertainties and other factors that may cause actual results with Afinitor to be materially different from any future results, performance or achievements expressed or implied by such statements. There can be no guarantee that Afinitor will be submitted or approved for any additional indications or labeling in any market. Nor can there be any guarantee that Afinitor will achieve any particular levels of revenue in the future. In particular, management's expectations regarding Afinitor could be affected by, among other things, unexpected regulatory actions or delays or government regulation generally; unexpected clinical trial results, including unexpected new clinical data and unexpected additional analysis of existing clinical data; the company's ability to obtain or maintain patent or other proprietary intellectual property protection; government, industry and general public pricing pressures; competition in general; the impact that the foregoing factors could have on the values attributed to the Novartis Group's assets and liabilities as recorded in the Group's consolidated balance sheet, and other risks and factors referred to in Novartis AG's current Form 20-F on file with the US Securities and Exchange Commission. Should one or more of these risks or uncertainties materialize, or should underlying assumptions prove incorrect, actual results may vary materially from those anticipated, believed, estimated or expected. Novartis is providing the information in this press release as of this date and does not undertake any obligation to update any forward-looking statements contained in this press release as a result of new information, future events or otherwise.

#### About Novartis

Located in East Hanover, New Jersey, Novartis Pharmaceuticals Corporation is an affiliate of Novartis AG, which provides healthcare solutions that address the evolving needs of patients and societies. Focused solely on healthcare, Novartis offers a diversified portfolio to best meet these needs: innovative medicines, eye care, cost-saving generic pharmaceuticals, consumer health products, preventive vaccines and diagnostic tools. Novartis is the only company with leading positions in these areas. In 2010, the Group's continuing operations achieved net sales of USD 50.6 billion, while approximately USD 9.1 billion (USD 8.1 billion excluding impairment and amortization charges) was invested in R&D throughout the Group. Headquartered in Basel, Switzerland, Novartis Group companies employ approximately 119,000 full-time-equivalent associates and operate in more than 140 countries around the world. For more information, please visit

<http://www.us.novartis.com>.

# Rapamune® (sirolimus) and Torisel® (temsirolimus) are registered trademarks of Wyeth Pharmaceuticals Inc.

## References

1. Yao, et al. Everolimus for Advanced Pancreatic Neuroendocrine Tumors. *New Eng J Med* 2011;364:514-23.
2. Halfdanarson, et al. Pancreatic neuroendocrine tumors (PNETs): incidence, prognosis and recent trend toward improved survival. *Annals of Onc* 19: 1727-1733, 2008.
3. National Library of Medicine and the National Institutes of Health. Pancreatic islet cell tumor. Available at <http://www.nlm.nih.gov/medlineplus/ency/article/000393.htm>. Accessed April 2011.
4. Novartis data on file. <http://www.pharma.us.novartis.com/product/pi/pdf/afinitor.pdf>. Accessed May 2011.
5. US Food and Drug Administration. Drug Details. ZANOSAR. Available at: <http://www.accessdata.fda.gov/scripts/cder/drugsatfda/index.cfm?fuseaction=Search.DrugDetails>. Accessed April 2011.
6. Yao, et al. One Hundred Years After "Carcinoid:" Epidemiology of and Prognostic Factors for Neuroendocrine Tumors in 35,825 Cases in the United States. *Journal of Clinical Oncology*. June 20 2009; vol. 26, number 18.
7. Motzer, et. al. Phase 3 Trial of Everolimus for Metastatic Renal Cell Carcinoma. *Cancer* 2010 Sep; 116(18):4256-4265.
8. National Library of Medicine and the National Institutes of Health. Neuroendocrine Tumor. Available at <http://www.cancer.gov/dictionary/?Cdrid=44904>. Accessed April 2011.
9. American Cancer Society Detailed Guides. Gastrointestinal Carcinoid Tumors. Available at <http://www.cancer.org/acs/groups/cid/documents/webcontent/003102-pdf.pdf>. Accessed April 2011.
10. American Cancer Society Detailed Guides. Pancreatic Cancer. Available at <http://www.cancer.org/acs/groups/cid/documents/webcontent/003131-pdf.pdf>. Accessed April 2011.

For Novartis multimedia content, please visit [www.thenewsmarket.com/Novartis](http://www.thenewsmarket.com/Novartis). For questions about the site or required registration, please contact: [journalisthelp@thenewsmarket.com](mailto:journalisthelp@thenewsmarket.com).

## Novartis Media Relations

Media only:

Investors only:

Denise Brashear

Richard Jarvis

Novartis Oncology

Novartis Corporation

P: +1 862 778 7336

P: +1 212 830 2433

Dana Kahn Cooper      [investor.relations@novartis.com](mailto:investor.relations@novartis.com)

P: +1 732 817 1800

F: +1 732 817 1834

SOURCE Novartis Pharmaceuticals Corporation

---

**Source URL:** <https://qa1.novartis.us/us-en/news/media-releases/novartis-gains-fda-approval-afinitor-first-new-treatment-nearly-three-decades-patients-advanced-pancreatic-net>

**List of links present in page**

1. <https://qa1.novartis.us/us-en/us-en/news/media-releases/novartis-gains-fda-approval-afinitor-first-new-treatment-nearly-three-decades-patients-advanced-pancreatic-net>
2. <http://www.afinitor.com/pNET>
3. <http://www.pharma.us.novartis.com/product/pi/pdf/afinitor.pdf>
4. <http://www.us.novartis.com/>
5. <http://www.nlm.nih.gov/medlineplus/ency/article/000393.htm>
6. <http://www.pharma.us.novartis.com/product/pi/pdf/afinitor.pdf>
7. <http://www.accessdata.fda.gov/scripts/cder/drugsatfda/index.cfm?fuseaction=Search.DrugDetails>
8. <http://www.cancer.gov/dictionary/?CdrID=44904>
9. <http://www.cancer.org/acs/groups/cid/documents/webcontent/003102-pdf.pdf>
10. <http://www.cancer.org/acs/groups/cid/documents/webcontent/003131-pdf.pdf>
11. <http://www.thenewsmarket.com/Novartis>
12. <mailto:journalisthelp@thenewsmarket.com>
13. <mailto:investor.relations@novartis.com>